

EFFECT OF POWDERED GREEN TEA MATCHA ON BIOFILM FORMATION BY MUTANS STREPTOCOCCI

Mochamad Fahlevi Rizal^{1,2a} , Noboru Kaneko^{1a*} , Hiroshi Ogawa^{1b} ¹Division of Preventive Dentistry, Department of Oral Health Science, Graduate School of Medical and Dental Sciences, Niigata University, Japan²Department of Pediatric Dentistry, Faculty of Dentistry, Universitas Indonesia, Indonesia^aDDS, PhD, Assistant Professor^bDDS, MDS, PhD, Professor and Head

ABSTRACT

DOI: [https://doi.org/10.25241/stomaeduj.2018.5\(4\).art.2](https://doi.org/10.25241/stomaeduj.2018.5(4).art.2)

Introduction: Antibacterial ingredients with high safety and mild taste that can be used for mouthwash for children are required. Matcha is one of the natural beverages that is made from the leaves of green tea (*Camellia sinensis*) and widely consumed by Japanese and Asian people. This study aimed to evaluate the effectiveness of matcha in suppressing biofilm formation by mutans streptococci, typical cariogenic bacteria.

Methodology: Five laboratory strains of *Streptococcus mutans* (serotype c, e and f) and *Streptococcus sobrinus* (serotype d and g) were used to evaluate the antibacterial effect of matcha. Matcha extract was added to bacterial cells in Heart Infusion broth supplemented with 1% sucrose (HIS). After incubation for 24 hours, the formed biofilm was dyed by Crystal Violet, and optical density at 490 nm was determined as the amount of biofilm formation. The effect of 0.02% chlorhexidine in HIS and purified water (MilliQ) in HIS were also measured as well.

Results: The amounts of biofilm formation by all serotypes of mutans streptococci in matcha + HIS were significantly lower than those in MilliQ + HIS as were chlorhexidine in HIS, except *S. sobrinus* serotype g. The differences of amounts biofilm formation by all *S. mutans* (serotype c, e and f) and *S. sobrinus* serotype d in matcha + HIS and in chlorhexidine + HIS were not significant statistically.

Conclusion: Matcha extract has an equivalent effectiveness of 0.02% chlorhexidine against most serotypes of mutans streptococci.

Keywords: *Streptococcus mutans*, *Streptococcus sobrinus*, chlorhexidine, green tea, mouthwash.

 OPEN ACCESS This is an Open Access article under the CC BY-NC 4.0 license.

 Peer-Reviewed Article

Citation: Rizal MF, Kaneko N, Ogawa H. Effect of powdered green tea matcha on biofilm formation by mutans Streptococci. Stoma Edu J. 2018;5(4):220-226.

Academic Editor: Mircea Ioan Popa MD, PhD, Professor, Head Microbiology Department II, "Carol Davila" University of Medicine and Pharmacy, Bucharest, Romania

Received: November 30, 2018
Revised: December 07, 2018
Accepted: December 17, 2018
Published: December 18, 2018

*Corresponding author: Assistant Professor Noboru Kaneko, DDS, PhD, Division of Preventive Dentistry, Department of Oral Health Science, Graduate School of Medical and Dental Sciences, Niigata University, 2-5274, Gakkocho-dori, Chuo-ku, Niigata City, Japan
Tel: +81 25 227 2857,
Fax: +81 25 227 0807,
e-mail: nkaneko@dent.niigata-u.ac.jp

Copyright: © 2018 the Editorial Council for the Stomatology Edu Journal.

1. Introduction

Efforts to obtain antibacterial ingredients that meet the criteria of comfort, especially taste, are needed to improve the health of the oral cavity. Oral health is inseparable from the normal flora balance of the oral cavity in order to prevent the occurrence of caries and gingivitis. The use of natural ingredients that are commonly consumed by the wider community in a place needs to be expanded in the functions of maintaining public health, especially oral health[1]. The ease of getting these and the consumption habits of the community makes it easy for materials like these to be accepted by the community if proven to have good effects on oral health. Because caries and gingivitis are also a problem in children in many countries, especially with developmental disabilities, an antibacterial made from natural ingredients for the oral cavity is urgently needed to be easily accepted by the child community, both in terms of safety and ease of use[2-4]. Each community has unique ingredients developed in its area, but has not being utilized them optimally in addition to regular consumption of food and beverages[5].

Green tea (*Camellia sinensis*) is commercially available in the market in three forms: Loose leaf, bagged and powdered (matcha)[6]. Matcha is generally consumed in Japan as a beverage that has high anti-oxidants such as flavonoids[7]. Many natural foods and beverages that are commonly consumed by people in a region, such as honey in the tropical regions, olive oil, virgin coconut oil and others have been partially known to have antibacterial effects especially in the oral cavity[8]. With many natural sources like ginger and garlic that also have flavonoid and are commonly used in Asia one needs to observe the antibacterial effect on oral bacteria[9]. As a comparison, chlorhexidine has been used for many years as a strong antibacterial standard and is also used in dentistry for preventive measures and treatment but the taste becomes an issue especially for children[10]. The streptococci that are commonly related to the human caries process are *Streptococcus mutans* and *Streptococcus sobrinus*. These cariogenic bacteria are divided into 5 serotypes (c, e, f, d and g)[11]. There has been no study on the antibacterial effects of matcha against each serotype of mutans streptococci. It

Table 1. Strains of mutans streptococci used in this study.

Strain	Serotype
<i>Streptococcus mutans</i>	
ATCC25175	c (type strain)
LM7	e
OMZ175	f
<i>Streptococcus sobrinus</i>	
ATCC33478	d (type strain)
6715	g

Table 2. Composition of cultures for biofilm assay.

	Antibacterial solution	Bacterial cells in HIS	HIS	MilliQ	Total
PC	0 µl	100µl	100µl	0 µl	200µl
MT	100µl	100µl	0 µl	0 µl	200µl
CHX	100µl	100µl	0 µl	0 µl	200µl
MQ	0µl	100µl	0 µl	100µl	200µl
NC	0 µl	0 µl	0 µl	200µl	200µl

PC: positive control; MT: matcha solution; CHX: 0.02% chlorhexidine gluconate; MQ: milliQ solution; NC: negative control

is expected that the results of research related to antibacterial from natural sources can be used to prevent infection in order to minimize chemical substances [12]. This study aimed to evaluate the effectiveness of matcha in suppressing biofilm formation by each serotype of mutans streptococci.

2. Material and Methods

2.1. Preparation of antibacterial solutions

As a matcha solution, 5 gram of powdered green tea matcha (Itoen, Tokyo, Japan) were added to 100 ml of water at 100°C and boiled for 15 minutes to extract. Prior to use in the laboratory, the solution was filtered using 0.20 µm Minisart filter (Sartorius, Göttingen, Germany). The aqueous chlorhexidine solution used in this study was 0.02% and was prepared from 20% chlorhexidine gluconate (Wako Pure Chemical, Osaka, Japan).

2.2. Culture of mutans streptococci

As mutans streptococci, we used 3 laboratory strains of *S. mutans* (serotype c, e, f) and 2 laboratory strain of *S. sobrinus* (serotype d, g) (table 1). Each strain from -80°C stock was cultured in Brain Heart Infusion (BHI) broth (Becton Dickinson, Maryland, USA) at 37°C overnight and then also sub-cultured in BHI broth. The bacterial solution was centrifuged and dissolved in Heart Infusion broth (Becton Dickinson, Maryland, USA) supplemented with 1% sucrose (HIS) until it reached 1×10^6 CFU/ml which was then used for the biofilm assay as strain culture.

2.3. Biofilm assay

The biofilm formation of each strain was evaluated using a 96 well plate. Each plate was divided into 5 areas of 12 wells and labeled as MT for wells that were

filled with matcha solution and strain culture, CHX for wells that were filled with 0.02% chlorhexidine and strain culture, MQ for wells that were filled with purified water (MilliQ) and strain culture, NC for wells that were filled with MilliQ alone as negative control and PC for wells that were filled with HIS and strain culture as positive control (Table 2). After 24 hours of incubation, the culture supernatant was discarded, and each well was washed twice with 200 µl Phosphate Buffered Saline (PBS). A 200 µl volume of 0.5% crystal violet was then added and incubated at 37°C for 15 minutes. The remaining crystal violet solution was discarded, the biofilm was washed once with 200 ml PBS, and then 200 µl of 96% ethanol was added to each well. The optical density of the biofilm formation was then measured at 490 nm (OD_{490}) using a microplate reader[13].

2.4. Statistical analysis

The OD_{490} values evaluated as biofilm formation were compared between NC, MQ and PC to confirm the validity of the measurement. Then OD_{490} values of MT, CHX, NC and MQ were compared to determine the differences in their effects on biofilm formation. In order to compare the differences, t-test and Bonferroni correction were used. All statistical analyses were performed using SPSS statistics 19 (IBM, USA); p values of 0.05 or less were considered significant.

3. Results

Fig. 1 shows OD_{490} values which mean amount of biofilm formation of NC (MilliQ alone), MQ (MilliQ + bacterial cells in HIS) and PC (bacterial cells in HIS) by *S. mutans* (serotype c, e and f) and *S. sobrinus* (serotype d and g), respectively. The OD_{490} of PC showed




Figure 1. Biofilm formation in NC (MilliQ alone), MQ (MilliQ + bacterial cells in HIS) and PC (bacterial cells in HIS) by each strain of mutans streptococci: (a) *S. mutans* ATCC25175 serotype c, (b) *S. mutans* LM7 serotype e, (c) *S. mutans* OMZ175 serotype f, (d) *S. sobrinus* ATCC33478 serotype d, (e) *S. sobrinus* 6715 serotype g.

significantly higher values than MQ, and those of MQ showed significantly higher values than NC in all strains (serotype c, e, f, d and g).

Fig. 2 shows OD₄₉₀ values which mean amount of biofilm formation of MT (matcha solution + bacterial cells in HIS), CHX (CHX solution + bacterial cells in HIS), NC (MilliQ alone) and MQ (MilliQ + bacterial cells in HIS), respectively. The OD₄₉₀ of MT, CHX and NC showed significantly lower values than MQ and no significant difference was found between MT, CHX and NC in all *S. mutans* (serotype c, e and f) and *S. sobrinus* (serotype d). In *S. sobrinus* serotype g, the OD₄₉₀ of MT showed lower value than MQ though the difference did not reach statistical significance. The OD₄₉₀ of CHX and NC showed significantly lower values than MQ and MT in *S. sobrinus* serotype

4. Discussion

The OD₄₉₀ of PC showed significantly higher values than MQ and those of MQ showed significantly higher values than NC in all strains, which indicated

that the biofilm assay used in this study could evaluate the amount of biofilm formation adequately. The OD₄₉₀ of MT and CHX showed significantly lower values than MQ and the differences between MT, CHX and NC were not significant in all strains except *S. sobrinus* serotype g. This means that the antibacterial effect of matcha solution resemble CHX solution in most strains of mutans streptococci.

This in-vitro research provides an overview of the antibacterial effects of matcha on various strains of mutans streptococci.

Mutans streptococci are considered as typical cariogenic bacteria[14,15]. Mutans streptococci can be classified into five serotypes, with dominant serotypes being slightly different locally. The effect of matcha against *S. sobrinus* serotype g was not significant among all mutans streptococci strains of serotype examined in this study. However, in the oral cavity of humans worldwide, *S. sobrinus* serotype g is not a type that is found predominantly[16-19]. In another study *S. sobrinus* serotype g showed




Figure 2. Effect of matcha extract and chlorhexidine on biofilm formation by each strain of mutans streptococci:

- (a) *S. mutans* ATCC25175 serotype c,
 (b) *S. mutans* LM7 serotype e,
 (c) *S. mutans* OMZ175 serotype f,
 (d) *S. sobrinus* ATCC33478 serotype d,
 (e) *S. sobrinus* 6715 serotype g.

more resistance to chlorhexidine than *S. mutans* and *S. sobrinus* serotype d[20]. Therefore, matcha is thought to have the effect of inhibiting biofilm formation for most of mutans streptococci in the oral cavity. One of the cariogenic properties of these bacteria is their high performance of glucan synthesis (extracellular polysaccharides) from sucrose, which contributes to the biofilm formation[21,22].

So, the effectiveness of matcha in suppressing the biofilm formation by mutans streptococci was important to be evaluated in this study.

A superior antibacterial would suppress more types of strains. However, caries is not only caused by a single bacterial species, but related to the biofilm activity involving many different bacteria species[23]. Chlorhexidine is used as a common antibacterial agent and is widely used for mouthwashes[24-28]. However, chlorhexidine has a taste issue, and it has problems such as side effects on the mucosa and it is staining the teeth, so it is not suitable for mouthwashes for children[29,30]. The concentration of

chlorhexidine used clinically is 0.12% and 0.2%, while in this study using a concentration of 0.02%. It was proved that the concentration of 0.03% was as effective as 0.12% or 0.2% concentration clinically[31-33]. The result of this study revealed that the low concentration of 0.02% chlorhexidine can inhibit biofilm formation of mutans streptococci.

Matcha has long been used for drinking and the safety has been established. The requirements for a mouthwash that can be used by children are related to safety issues[34,35]. It is also considered to have less problems with taste. Therefore, matcha can be considered a suitable ingredient for mouthwashes for children.

The bacteria investigated this time are mutans streptococci and are strongly related to dental caries. However, there are many other bacteria in the oral cavity related to dental caries and other oral diseases such as periodontitis. In this study, we did not consider the influence on other such cariogenic bacteria and on periodontal disease causing bacteria. Hereafter,

it seemed necessary to investigate the antibacterial effect on those bacteria as well.

5. Conclusion

Matcha has the equivalent effectiveness of 0.02% chlorhexidine against most serotypes of mutans streptococci, namely *S. mutans* serotype c, e and f and *S. sobrinus* serotype d. It is suggested that matcha is a promising ingredient as a safe and effective for mouthwashes in children.

Author Contributions

MFR: Idea, experimental design, data analysis, wrote the manuscript. NK: Performed biofilm and spectro-

photometry assay, substantial contributed to writing manuscript. HO: Performed substantial contributed to writing manuscript.

Acknowledgement

The authors have no conflict of interest. This research funding by Division of Preventive Dentistry, Department of Oral Health Science, Graduate School of Medical and Dental Sciences, Niigata University.

The authors declare that the research was conducted in the absence of any commercial or financial relationship that could be construed as a potential conflict of interest.

References

- Jacob, PS. Editorial: Natural products in oral health *The Open Dent J.* 2016;10(Suppl-1, M1):158-159. doi: 10.2174/1874210601610010158. eCollection 2016.
- Norwood KW, Slayton, RL. Oral health care for children with developmental disabilities. *Pediatrics.* 2013;131(3):614-619. doi: 10.1542/peds.2012-3650. [Full text links] [PubMed] Google Scholar(53) Scopus(22)
- Lachenmeier DW, Monokhova YB, Markova M, et al. What happens if people start drinking mouthwash as surrogate alcohol? A quantitative risk assessment. *Food Chem Toxicol.* 2013 Jan; 51:173-178. doi: 10.1016/j.fct.2012.09.031. [Full text links] [PubMed] Google Scholar(13) Scopus(8)
- Werner CWdeA, Seymour RA. Are alcohol containing mouthwashes safe? *Br Dent J.* 2009;207(10): E19; discussion 488-489. doi: 10.1038/sj.bdj.2009.1014. [Full text links] [PubMed] Google Scholar(37) Scopus(16)
- Palombo EA. Traditional medicinal plant extracts and natural products with activity against oral bacteria: potential application in the prevention and treatment of oral diseases. *Evid Based Complement Alternat Med.* 2011;2011:680354. doi: 10.1093/ecam/nep067. [Full text links] [Free PMC Article] [PubMed] Google Scholar(381) Scopus(196)
- Farooq S, Sehgal A. Antioxidant activity of different forms of green tea: loose leaf, bagged and matcha. *Curr Res Nutr Food Sci.* 2018;6(1):35-40. doi : <http://dx.doi.org/10.12944/CRNFSJ.6.1.04> [Full text links] Google Scholar(1)
- Dietz C, Dekker M, Fiszman BP. An intervention study on the effect of matcha tea, in drink and snack bar formats, on mood and cognitive performance. *Food Res Int.* 2017; 99(Pt 1):72-83. doi: 10.1016/j.foodres.2017.05.002. [Full text links] [PubMed] Google Scholar(5) Scopus(3)
- Lavine P, Fauziah E, Rizal MF, Budiardjo SB. Antibacterial effect of virgin coconut oil on (*Actinomyces* Sp.) that causes dental black stain in children. *Asian J Pharm Clin Res.* 2018;11(2): 333-335. doi: <http://dx.doi.org/10.22159/ajpcr.2018.v11i2.23199>
- Ali AMA, El-Nour MEM, Yagi SM. Total phenolic and flavonoid contents and antioxidant activity of ginger (*Zingiber officinale* Rosc.) rhizome, callus and callus treated with some elicitors. *J Genet Eng Biotechnol.* 2018. <https://doi.org/10.1016/j.jgeb.2018.03.003> [Full text links] Google Scholar(2) Scopus(1)
- Calogiuri GF, Di Leo E, Trautmann A, et al. Chlorhexidine hypersensitivity: a critical and updated review. *J Allergy Ther.* 2013;4(4):1-7. doi: 10.4172/2155-6121.1000141 [Full text links] Google Scholar(35)
- Coykendall AL, Bratthall D, O'Connor K, Dvarskas RA. Serological and genetic examination of some nontypical *Streptococcus mutans* strains. *Infect Immun.* 1976;14(3):667-670. [Full text links] [Free PMC Article] [PubMed] Google Scholar(48) Scopus(17)
- Kiany F, Niknahad H, Niknahad M. Assessing the effect of pomegranate fruit seed extract mouthwash on dental plaque and gingival inflammation. *J Dent Res Rev.* 2016;3(4):117-123. doi: 10.4103/2348-2915.200016 [Full text links] Google Scholar(1)
- Linggriani A, Rizal MF, Fauziah E, Suharsini M. Differences in the effect of 0.05% and 0.1% propolis flavonoids on in vitro biofilm formation by *Streptococcus mutans* from children's dental plaque. *Asian J Pharm Clin Res.* 2018;11(2):215-218. doi: <http://dx.doi.org/10.22159/ajpcr.2018.v11i2.22521>
- Sánchez-Acedo M, Montiel-Company JM, Dasi-Fernández F, Almerich-Silla JM. *Streptococcus mutans* and *Streptococcus sobrinus* detection by Polymerase Chain Reaction and their relation to dental caries in 12 and 15 year-old schoolchildren in Valencia (Spain). *Med Oral Patol Oral Cir Bucal.* 2013;18(6): e839-e845. doi: 10.4317/medoral.18941 [Full text links] [Free PMC Article] [PubMed] Google Scholar(12) Scopus(12)
- Okada M, Kawamura M, Oda Y, et al. Caries prevalence associated with *Streptococcus mutans* and *Streptococcus sobrinus* in Japanese schoolchildren. *Int J Paediatr Dent.* 2012;22(5):342-348. doi: 10.1111/j.1365-263X.2011.01203.x. [Full text links] [PubMed] Google Scholar(31) Scopus(13)
- Nakano K, Nemoto H, Nomura R, et al. Serotype distribution of *Streptococcus mutans* a pathogen of dental caries in cardiovascular specimens from Japanese patients. *J Med Microbiol.* 2007;56(Pt4):551-556. doi: 10.1099/jmm.0.47051-0 [Full text links] [PubMed] Google Scholar(59) Scopus(38)
- Holbrook WP, Beighton D. *Streptococcus mutans* levels in saliva and distribution of serotypes among 9-year-old Icelandic children. *Scand J Dent Res.* 1987;95(1):37-42. [Full text links] [PubMed] Google Scholar(46)
- Fauziah E, Sutadi H, Bachtiar EW. Prediction of baby bottle tooth decay syndrome using the salivary mucin MG2. *Int J Clin Prev Dent.* 2013;9(4):211-214. [Full text links] Google Scholar(1)
- Kozai K, Nakayama R, Tedjosongko U, et al. Intrafamilial distribution of mutans streptococci in Japanese families and possibility of father-to-child transmission. *Microbiol Immunol.* 1999;43(2):99-106. doi: <https://doi.org/10.1111/j.1348-0421.1999.tb02380.x> [Full text links] [PubMed] Google Scholar(131) Scopus(57)
- Grönroos L, Mättö J, Saarela M, et al. Chlorhexidine susceptibilities of mutans streptococcal serotype and ribotypes. *Antimicrob Agents Chemother.* 1995;39(4):894-898. doi: 10.1128/AAC.39.4.894 [Full text links] [Free PMC Article] [PubMed] Google Scholar(51) Scopus(33)
- Hahnel S, Mühlbauer G, Hoffmann J, et al. *Streptococcus mutans* and *Streptococcus sobrinus* biofilm formation and metabolic activity on dental materials. *Acta Odontol Scand.* 2012;70(2):114-121. doi:

- 10.3109/00016357.2011.600703.
[Full text links] [PubMed] Google Scholar(13) [Scopus\(9\)](#)
22. Merritt J, Qi F. The mutacins of *Streptococcus mutans*: regulation and ecology. *Mol Oral Microbiol*. 2012;27(2):57-69. doi: 10.1111/j.2041-1014.2011.00634.x.
[Full text links] [Free PMC Article] [PubMed] Google Scholar(81) [Scopus\(59\)](#)
23. Sanz M, Beighton D, Curtis MA, et al. Role of microbial biofilms in the maintenance of oral health and in the development of dental caries and periodontal diseases. Consensus report of group 1 of the Joint EFP/ORCA workshop on the boundaries between caries and periodontal disease. *J Clin Periodontol*. 2017 Mar;44 Suppl 18:S5-S11. doi: 10.1111/jcpe.12682.
[Full text links] Google Scholar(45) [Scopus\(27\)](#)
24. Kocak MM, Ozcan S, Kocak S, et al. Comparison of the efficacy of three different mouthrinse solutions in decreasing the level of streptococcus mutans in saliva. *Eur J Dent*. 2009;3(1):57-61.
[Full text links] [Free PMC Article] [PubMed] Google Scholar(51)
25. Simratvir M, Singh N, Chopra S, Thomas AM. Efficacy of 10% povidone iodine in children affected with early childhood caries: an in vivo study. *J Clin Pediatr Dent*. 2010;34(3):233-238.
[Full text links] [PubMed] Google Scholar(30) [Scopus\(17\)](#)
26. Cieplik F, Kara E, Muehler D, et al. Antimicrobial efficacy of alternative compounds for use in oral care toward biofilms from caries-associated bacteria in vitro. *MicrobiologyOpen*. 2018 Jul 26:e00695. doi: 10.1002/mbo3.695.
27. Vinod KS, Sunil KS, Sethi P, et al. A novel herbal formulation versus chlorhexidine mouthwash in efficacy against oral microflora. *J Int Soc Prev Community Dent*. 2018;8(2):184-190. doi: 10.4103/jispcd.JISPCD_59_18
28. James P, Worthington HV, Parnell C, et al. Chlorhexidine mouthrinse as an adjunctive treatment for gingival health. *Cochrane Database Syst Rev*. 2017 Mar 31;3:CD008676. doi: 10.1002/14651858.CD008676.pub2.
[Full text links] [PubMed] Google Scholar(3) [Scopus\(23\)](#)
29. Gunasekaran G, Lakshmanan R. Effect of chlorhexidine mouthwash on taste alteration. *Asian J Pharm Clin Res*. 2016;9(13):102-104.
[Full text links] Google Scholar(2)
30. Bagis B, Baltacioglu E, Özcan M, Ustaomer S. Evaluation of chlorhexidine gluconate mouthrinse-induced staining using a digital colorimeter: an in vivo study. *Quintessence Int*. 2011;42(3):213-223.
[Full text links] [PubMed] Google Scholar(36) [Scopus\(18\)](#)
31. Bonesvoll P. Oral pharmacology of chlorhexidine. *J Clin Periodontol*. 1977;4(5):49-65.
[Free text links] [PubMed] Google Scholar(77) [Scopus\(46\)](#)
32. Dogan S, Günay H, Leyhausen G, Geurtsen W. Effects of low-concentrated chlorhexidine on growth of *Streptococcus sobrinus* and primary human gingival fibroblasts. *Clin Oral Investig*. 2003;7(4): 212-216. doi: 10.1007/s00784-003-0226-3.
[Free text links] [PubMed] Google Scholar(10) [Scopus\(8\)](#)
33. Sylvie S, Catherine F, Jean-Philippe G, Christine R. In vitro bactericidal assay under simulated practical conditions for comparison of chlorhexidine mouthrinses: chlorhexidine concentration is only one of the in vitro activity criteria. *Oral Health Dentl Manag*. 2016;15(6):1-6.
[Full text links] Google Scholar(0) [Scopus\(0\)](#)
34. Massey CC, Shulman JD. Acute ethanol toxicity from ingesting mouthwash in children younger than age 6, 1989-2003. *Pediatr Dent*. 2006;28(5):405-409. [Full text links] [PubMed] Google Scholar(12)
35. Sharma R, Hebbal M, Ankola AV, et al. Effect of two herbal mouthwashes on gingival health of school children. *J Tradit Complement Med*. 2014;4(4):272-278. doi: 10.4103/2225-4110.131373.
[Full text links] [Free PMC Article] [PubMed] Google Scholar(12) [Scopus\(6\)](#)

Mochamad Fahlevi RIZAL

DDS, PhD, Assistant Professor
Department of Oral Health Science
Division of Preventive Dentistry
Graduate School of Medical and Dental Sciences
Niigata University, Japan, Niigata, Japan



Mochamad Fahlevi Rizal is a Pediatric Dentist in Indonesia. Join become lecturer in Department of Pediatric Dentistry, Faculty of Dentistry, Universitas Indonesia since 1998. In 2018 become visiting lecturer in Niigata University and made collaborative research with Division of Preventive Dentistry, Department of Oral Health Science, Graduate School of Medical and Dental Sciences, Niigata University.

Questions

1. Which was the matcha used in this study?

- a. 5 grams in 100 ml water
- b. 0.5 gram in 100 ml water
- c. 0.05 gram in 100 ml water
- d. 50 gram in 100 ml water

2. What species of mutans streptococci were used in this study?

- a. *Streptococcus ferus* and *Streptococcus pneumoniae*
- b. *Streptococcus mutans* and *Streptococcus sobrinus*
- c. *Streptococcus macacae* and *Streptococcus rattii*
- d. *Streptococcus criceti* and *Streptococcus downei*

3. What was the antibacterial used to compare the effectiveness of matcha in this study?

- a. Ethil alcohol 70%;
- b. Povidone iodine;
- c. Chlorhexidine;
- d. Gentian violet.

4. What was the result of this study?

- a. All strain of mutans streptococci were suppressed significantly by matcha;
- b. Only serotype c was suppressed significantly by matcha;
- c. Only serotype c, e, and d were suppressed significantly by matcha;
- d. Only serotype c, e, f, and d were suppressed;



<http://www.efcd.eu/conseuro-meetings/>