

STABILITY OF SKELETAL CLASS III MALOCCLUSION AFTER ORTHOGNATHIC SURGERY AND ORTHODONTIC TREATMENT: A SYSTEMATIC REVIEW AND META-ANALYSIS

Abdolreza Jamilian^{1a*}, Ludovica Nucci^{2b}, Ali Fateh^{1c}, Mitra Toliat^{1d}, Alireza Darnahal^{3e}, Madi Alassadi^{3f}, Chin Wei Wang^{3g}

¹Department of Orthodontics, Cranio-Maxillofacial Research Center, Tehran University of Medical Sciences, Islamic Azad University, Tehran, Iran

²Multidisciplinary Department of Medical-Surgical and Dental Specialties, Dental School, University of Campania Luigi Vanvitelli, Naples, Italy

³Department of Periodontics and Oral Medicine, School of Dentistry, University of Michigan, Ann Arbor, MI 48109, USA

^aDDS, MSc, OMFS, Professor; e-mail: info@jamilian.net; ORCIDiD: <https://orcid.org/0000-0002-8841-0447>

^bUndergraduate Dental Student; e-mail: ludovica.nucci@virgilio.it; ORCIDiD: <https://orcid.org/0000-0002-7174-7596>

^cDDS, Executive Manager; e-mail: Af_fateh@yahoo.com; ORCIDiD: <https://orcid.org/0000-0002-1895-0632>

^dDDS; e-mail: Toliatmitra@gmail.com

^ePost Doctorate Student; e-mail: alirezadarnahal@gmail.com; ORCIDiD: <https://orcid.org/0000-0001-9629-731X>

^fPost Doctorate Student; e-mail: alassadi@umich.edu

^gDDS, Clinical Assistant Professor, Director of Predoctoral Periodontics; e-mail: Jeffwa@umich.edu; ORCIDiD: <https://orcid.org/0000-0003-2274-8901>

ABSTRACT

[https://doi.org/10.25241/stomaedu.2020.7\(1\).art.7](https://doi.org/10.25241/stomaedu.2020.7(1).art.7)

Background Relapse is one of the major concerns in the correction of skeletal class III malocclusion.

Objective The purpose of this systemic review was to evaluate the degree of relapse on skeletal class III patients who received bimaxillary surgery or mandibular setback with orthodontic treatment.

Data Sources A search of the literature was performed in the databases of PubMed, Google Scholar Beta, Scopus, Web of Science, and the Cochrane Library.

Study Selection Out of the 165 articles identified, 73 studies were obtained, once duplicated articles were excluded. Then, 40 other records were excluded due to titles and abstracts, and 20 were removed for not fulfilling exclusion/inclusion criteria. 11 studies met the final inclusion criteria. Some cephalometric data during T1–T2–T3 were measured.

Data Extraction SNA did not have any significant changes within less than 2 years but it increased significantly after 2 years. SNB did not have any significant changes in more than 2 years' follow-up, while it rose significantly in less than 2 years. Overjet decreased significantly after 2 years but not earlier than this duration. Overbite intensified significantly in more than 2 years and not earlier.

Data Synthesis SNA and overbite increased significantly after 2 years. SNB increased significantly before 2 years and did not have any changes after it. Overjet was significantly reduced after 2 years.

KEYWORDS

Class III; Skeletal and Dental Changes; Stability; Bimaxillary Surgery or Mandibular Setback; Systematic Review and Meta-Analysis.

1. INTRODUCTION

Moderate to severe skeletal class III patients often require a combined orthodontic and surgical approach for treatment. It has been reported that skeletal class III malocclusion is the most frequent deformity corrected by combined orthognathic

surgery and orthodontic treatment [1-4]. However, bimaxillary surgery has gradually become more popular to correct class III malocclusion [5-7]. It has been estimated that 20% to 25% of all Class III cases have true mandibular prognathism suggesting that at least 75% of all class III malocclusions have some degree of maxillary retrusion. Given this



OPEN ACCESS This is an Open Access article under the CC BY-NC 4.0 license.

Peer-Reviewed Article

Citation: Jamilian A, Nucci L, Fateh A, Toliat M, Darnahal A, Alassadi M, Wang CW. Stability of skeletal Class III malocclusion after orthognathic surgery and orthodontic treatment: a systematic review and meta-analysis. Stoma Edu J. 2020;7(1):52-67.

Received: February 06, 2020; **Revised:** February 13, 2020; **Accepted:** February 14, 2020; **Published:** February 17, 2020

***Corresponding author:** Professor Abdolreza Jamilian, Department of Orthodontics, TUMS School of Dentistry, Cranio-Maxillofacial Research Center, Tehran University of Medical Sciences, Islamic Azad University, Tehran, Iran, No 14, Pesiyan Ave., Vali Asr St, Tehran, 1986944768, Iran
Tel/Fax: 00982122011892, e-mail: info@jamilian.net

Copyright: © 2020 the Editorial Council for the Stomatology Edu Journal.




Figure 1. PRISMA flowchart of article retrieval.

scenario, the surgical treatment has been regarded as the best approach to achieve the best results in adult cases [8]. Post-surgical relapse is one of the major concerns in the correction of skeletal class III malocclusion. It has been shown that there is a greater tendency for relapse after bimaxillary osteotomy [9]. Similarly, LaBanc et al [10] reported that due to increased incidence of relapse, bimaxillary surgery should only be performed for specific indications; for example two-jaw surgery has greater relapse than single-jaw surgery. On the contrary, Proffit et al [11] found that better stability and predictable results can be obtained after two-jaw surgery. Immediate relapse can be identified after surgery which may occur due to intraoperative error such as imprecise planning, inaccurate osteosynthesis, or failure to fix the joint. On the other hand, late relapse can be detected once a considerable period has elapsed since the day of the surgical procedure. Late relapse may occur due to unstable occlusal relationships, growth spurts, absence of myofunctional adaptation, or persistent tongue or orofacial muscle habits [12]. A study [5] revealed that the rate of short-term

relapse of bilateral sagittal split osteotomy setback surgery is 9.9%-62.1%, and long-term relapse is between 14.9% and 28.0%, at point B. To the best of our knowledge, there has been no meta-analysis evaluating the stability of skeletal changes after a combined orthodontic and surgical procedure for treatment of skeletal Class III patients in short- and long-term follow ups. The question is: "How much of the treatment effects remains stable by the end of follow-up?"

Objectives: The aim of this study was a meta-analysis of the literature on the stability of skeletal class III malocclusion.

Patients often require a combined orthodontic and surgical approach after bimaxillary surgery. This meta-analysis was undertaken to explore the parameters related to the skeletal stability in surgical correction of skeletal Class III malocclusion.

2. MATERIALS AND METHODS

2.1. Protocol and registration

This systematic review was based on a specific protocol developed and piloted following the




Figure 2a. Before surgery T1 - After surgery T2, (T1-T2) SNA.




Figure 2b. After surgery T2 - Last follow-up T3, (T2-T3) SNA




Figure 2c. Subgroup analysis according to the follow-up period (SNA).

1. Less than 2 years.; 2. More than 2 years.




Figure 3a. Before surgery T1 - After surgery T2, (T1-T2) SNB.




Figure 3b. After surgery T2 - Last follow-up T3, (T2-T3) SNB.




Figure 3c. Subgroup analysis according to the follow-up period (SNB).

1. Less than 2 years.; 2. More than 2 years.




Figure 4a. Before surgery T1 - After surgery T2, (T1-T2) ANB.




Figure 4b. After surgery T2 - Last follow-up T3, (T2-T3) ANB.




Figure 4c. Subgroup analysis according to the follow-up period (ANB).

1. Less than 2 years ; 2. More than 2 years.




Figure 5a. Before surgery T1 - After surgery T2, (T1-T2) Overjet.




Figure 5b. After surgery T2 - Last follow-up T3, (T2-T3) Overjet.




Figure 5c. Subgroup analysis according to the follow-up period (Overjet).

1. Less than 2 years.; 2. More than 2 years.

Table 1. Eligibility criteria used for the study selection.

Category	Inclusion criteria	Exclusion criteria
Study design	Randomized controlled trials Controlled clinical trials Cohort studies Experimental studies, prospective and retrospective studies comparing at least two surgical treatment strategies Articles written in the English language	Case reports Commentaries Systematic reviews or meta-analyses descriptive studies, opinion articles, or abstracts
Participants	Non-growing patients with Class III malocclusion Follow-up of 6 months or longer	Patients with cleft lip palate and/or craniofacial syndromes or genetic problems Patients with temporomandibular joint disorders Treatment in growing patients
Intervention	Mandibular set back Bimaxillary surgery: (including maxillary advancement and mandibular set back) Surgery first	Maxillary advancement Patients treated with orthodontic or orthopedic appliances Studies not concerning surgical long-term stability
Outcome	Skeletal and dentoalveolar variables measured by lateral cephalometric radiographs	Studies providing no cephalometric measurements
Average time of follow-up	Studies with an average follow-up at least 2 years after surgery	Studies with no follow-up




Figure 7a. Before surgery T1 - After surgery T2, (T1-T2) U1/SN.




Figure 7b. After surgery T2 - Last follow-up T3, (T2-T3) U1/SN.




Figure 8a. Before surgery T1 - After surgery T2, (T1-T2) L1/MP.




Figure 8b. After surgery T2 - Last follow-up T3, (T2-T3) L1/MP.

guidelines outlined in the PRISMA-P statement [13]. Furthermore, the procedure and reporting followed the guidelines of the Cochrane Handbook for Systematic Reviews of Interventions [14] and the PRISMA statement, [15] respectively.

2.2. Information sources, search strategy, and study selection

A literature search was performed using PubMed, Google Scholar Beta, Scopus, Web of Science, and the Cochrane Library to identify articles reporting combined orthodontic and surgical approach for treatment of skeletal class III malocclusion in non-growing patients. The search process was conducted independently by two coauthors (AJ and AD) for articles published up to December 2018. All titles and abstracts were evaluated, and duplicate studies were removed.

2.3. Eligibility criteria

The inclusion and exclusion criteria were established prior to the search according to Table 1. Each keyword

was carefully selected and revised for each database. All keywords used in the search are detailed in Table 2. This systematic review and meta-analysis was conducted based on the PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) guidelines [16]. Title-abstract-full text of each article was checked independently by two coauthors based on the PRISMA chart.

2.4. Data collection and data items

Two authors (AJ and AD) used pre-defined electronic sheets to extract study characteristics independently. Three time points were defined: T1 (before surgery), T2 (after surgery), and T3 (the end of the follow-up). The findings were obtained on the following items: Name of first author, year of publication, country, number and mean age of patients, gender, type of surgery (Mandibular setback including BSSO or vertical osteotomy) or (Bimaxillary surgery including Lefort 1 + BSSO or Lefort 1 + vertical), type of fixation including Rigid Internal Fixation (RIF) or Maxillomandibular

Table 2. Keywords used for each data base search.

Pubmed	Web of science	Scopus	Embase	Cochrane
50	7	63	42	3
(((((("Malocclusion, Angle Class III"[Mesh] OR "class3"[Title/Abstract]) OR "class III"[Title/Abstract]) OR "Maxillary Deficiency"[Title/Abstract]) OR "mandibular protrusion"[Title/Abstract]) OR "Maxillary retrusion"[Title/Abstract]) AND (((("Mandibular Osteotomy"[Mesh] OR "mandibular surgery"[Title/Abstract]) OR "bimaxillary surgery"[Title/Abstract]) OR "surgical orthodontics"[Title/Abstract]) OR "mandibular set back"[Title/Abstract])) AND (stability[Title/Abstract] OR relapse[Title/Abstract]))	TI= ("Malocclusion, Angle Class III" OR "class3" OR "class III" OR "Maxillary Deficiency" OR "mandibular protrusion" OR "Maxillary retrusion") AND TI= ("Mandibular Osteotomy" OR "mandibular surgery" OR "bimaxillary surgery" OR "surgical orthodontics" OR "mandibular set back") AND TI= (stability OR relapse)	((TITLE-ABS-KEY ("Malocclusion, Angle Class III") OR TITLE-ABS-KEY ("class3") OR TITLE-ABS-KEY ("class III") OR TITLE-ABS-KEY ("Maxillary Deficiency") OR TITLE-ABS-KEY ("mandibular protrusion") OR TITLE-ABS-KEY ("Maxillary retrusion")) AND ((TITLE-ABS-KEY ("Mandibular Osteotomy") OR TITLE-ABS-KEY ("mandibular surgery") OR TITLE-ABS-KEY ("bimaxillary surgery") OR TITLE-ABS-KEY ("surgical orthodontics") OR TITLE-ABS-KEY ("mandibular set back"))) AND ((TITLE-ABS-KEY (stability) OR TITLE-ABS-KEY (relapse))))	'malocclusion angle class iii':ab,ti OR 'class3':ab,ti OR 'class III':ab,ti OR 'maxillary deficiency':ab,ti OR 'maxillary retrusion':ab,ti and 'mandibular osteotomy':ab,ti OR 'mandibular surgery':ab,ti OR 'bimaxillary surgery':ab,ti OR 'surgical orthodontics':ab,ti OR 'mandibular set back':ab,ti and 'stability':ab,ti OR 'relapse':ab,ti	("Malocclusion, Angle Class III" OR "class3" OR "class III" OR "Maxillary Deficiency" OR "mandibular protrusion" OR "Maxillary retrusion") AND ("Mandibular Osteotomy" OR "mandibular surgery" OR "bimaxillary surgery" OR "surgical orthodontics" OR "mandibular set back") AND (stability OR relapse)

Fixation (MMF), follow-up after surgery until 2 years, follow-up after surgery until 5 years, SNA, SNB, ANB, overjet, overbite, incisor mandibular plane angle (IMPA), upper incisor to SN (U1/SN) angle, during T1–T2 (surgical effects), T2–T3 (posttreatment changes) were recorded. In order to identify the correlation between relapse and cephalometric landmarks, meta-analyses were conducted between cephalometric landmarks and different variables such as type of surgery (mandibular setback including BSSO or vertical osteotomy) or (bimaxillary surgery including Lefort 1 + BSSO or Lefort 1 + vertical), type of fixation including Rigid Internal Fixation (RIF) or Maxillomandibular Fixation (MMF), follow-up after surgery within 2 years, follow-up after surgery within 5 years. The cut-off value of less than 2 years was chosen to separate short-term from long-term studies.

2.5. Statistical analysis

All statistical tests were conducted using the STATA 14 (StataCorp LP, College Station, USA). The effects of bimaxillary surgery or mandibular setback on SNA, SNB, ANB, overjet, overbite, incisor mandibular plane angle (IMPA) and upper incisor to SN (U1/SN) angle, before and after surgery as well as the last follow-up were measured by weighted mean difference (WMD) and the 95% confidence intervals (CI). The standard error (SE) of the mean difference (MD) for non-reported studies was calculated by the following formula: $SD^2 \text{ baseline} + SD^2 \text{ final} - (2 R^* SD \text{ baseline} + SD \text{ final})$ and $SD = SE * \sqrt{n}$. Heterogeneity across studies was assessed using the I-squared and the alpha of 0.05 for statistical significance. The subgroup analysis was based on the time of

follow-up to identify the source of heterogeneities. To identify the source of clinical heterogeneity, susceptible variables including treatment plan, gender, country, treatment time, type of surgery, type of fixation, and follow-up time were introduced into a meta-regression model. WMD with 95% CI was calculated for all variables. The publication bias was determined using Begg tests. The p-value of 0.05 was regarded for statistical significance. The changes in seven variables (SNA, SNB, ANB, IMPA, overjet, U1/SN, and L1/MP) during three time periods [Before surgery (T1), after surgery (T2) and last follow-up (T3)] were compared between the studies. The summarized data of included studies and cephalometric measurements of the included studies are seen in Tables 3 and 4 respectively. The results of the statistical analysis for heterogeneity and the funnel plots are displayed in Figures 2, 3, 4, 5, 6, 7, and 8.

3. RESULTS

3.1. Study selection and characteristics

Of the 165 records resulting from the search strategies, 73 studies were obtained once duplicated articles were excluded. Then, 40 papers were removed because of their titles and abstracts. In addition, 20 more were further excluded for not meeting the exclusion/inclusion criteria. Ultimately, 13 papers met the final selected criteria and were selected to conduct the systematic review and meta-analysis. The manual search did not yield any additional material. In case of disagreement, the authors discussed the controversy until an agreement was reached. Of the 13 studies, 2 of

Table 3. Characteristics of the included studies in the systematic review.

Publication	Observation period	Group(s)	Participants (Number, sex, age)	Type of Surgery	Type of fixation	Country
Jeong et al [1], 2018	2 years	conventional bimaxillary surgery surgery first	14 (7M, 7F) / Mean age (21.5 ± 2.5) 17 (9M, 8F) / Mean age (20.3 ± 2.2)	Le Fort I Osteotomy & Bilateral IVRO*	Rigid Wire	South Korea
Choi et al [2], 2016	2 years	IVRO IVRO + Lefort 1	15 (7M, 8F) / Mean age (22.1 ± 2.3) 15 (7M, 8F) / Mean age (24.1 ± 4.3)	IVRO IVRO + Lefort 1	Intermaxillary Fixation Rigid	South Korea
Park et al [3], 2016	6 months	Conventional bimaxillary surgery Surgery-first	20 (13M, 7F) / Mean age (25.3) 20 (12M, 8F) / Mean age (22.6)	Lefort 1 & BSSO**	Rigid	South Korea
Aydemir et al [4], 2015	5 years	BSSO with Semi Rigid Fixation and Lefort 1 with Rigid Fixation	26 / Mean age (17-29)	Lefort 1 & BSSO**	Semi Rigid and Rigid Fixation	Turkey
Kim et al [5], 2014	1 year	Bimaxillary Surgery Le Fort 1 and IVRO	37 (20M, 17F) / Mean age (23 ± 4)	Lefort 1 & IVRO	Wire	South Korea
Kim et al [6], 2014	22 months	Conventional-BSSO Surgery first	23 (14M, 9F) / Mean age (23 ± 6.3) 38 (19M, 19F) / Mean age (21.6 ± 3.5)	SF*** BSSO	Rigid Rigid	South Korea
Ko et al [7], 2013	1 year		45 (19M, 26F) / Mean age (23.2)	Lefort I & bilateral sagittal split osteotomy	Monocortical plates and screws in each side of the mandible and miniplates in each side of the maxilla	Taiwan
Jakobsone et al [8], 2011	3 years	Lefort 1 + BSSO	81 (55M, 26F) / Mean age (25.8 ± 9.5)	Lefort 1 + BSSO	Rigid	Norway
Abeltins et al [9], 2011	1 year	IVRO BSSO	21 (6M, 15F) / Mean age (20.2) 30 (7M, 23F) / Mean age (19.4)	Lefort 1 + BSSO Lefort 1 + IVRO	Wire	Latvia
Iannetti et al [10], 2007	2 years	Lefort 1 Lefort 1 + BSSO	N/A / Mean age (18-36)	Bimaxillary Lefort 1	Rigid	Italy
Costa et al [11], 2006	1 year	rigid titanium rigid resorbable plate	21 / Mean age (N/A)	Rigid resorbable plate Rigid titanium	Rigid resorbable plate Rigid titanium	Italy
Kwon et al [12], 2000	7 months	Lefort 1 and BSSO	25 (13M, 12F) / Mean age (24.1)	Lefort 1 + BSSO	Rigid	Japan
McCance et al [13], 1992	1 year	Lefort 1 + BSSO	18 (N/A) / Mean age (N/A)	Lefort 1 + BSSO	Wire	England

Age is reported in years; N/A, Not available; M, Male; F, Female

*IVRO, Intra-oral vertical ramus osteotomy; **BSSO, Bilateral sagittal split osteotomy; SF

them did not provide us with enough data for the meta-analysis and were excluded from the study. Performing meta-analysis was only feasible for 11 studies, and these studies were included in our study. These studies include non-growing patients with Class III malocclusion with Follow-up of 6 months or longer. The level of inter-examiner agreement of data extraction was measured using kappa statistics. The level of agreement between the two examiners was assessed using the Cohen kappa scores. The kappa score for study selection was 0.978, indicating an excellent level of agreement. The PRISMA flow diagram of study selection is outlined in Fig. 1. Before surgery T1– After surgery T2, (T1–T2) Figures 2a, 3a, 4a, 5a, 6a, 7a, and 8a.

Short-term treatment effects included significant increase in SNA (WMD 1.78, 95%CI:1.42, 2.12), significant reduction in SNB (WMD -3.95, 95%CI:-4.50, -3.40), significant rise in ANB (WMD 6.36, 95%CI:5.59, 7.13), significant growth in overjet (WMD 8.24, 95%CI: 6.56, 9.92), significant elevation in overbite (WMD 1.57, 95%CI: 0.30,2.84), while U1/SN (WMD -2.34, 95%CI, -6.27, 1.58) and L1/MP (WMD 2.12, 95%CI:0.43, 3.82) did not show any significant changes. After surgery T2– Last follow upT3, (T2–T3) Figures 2b, 3b, 4b, 5b, 6b, 7b, and 8b.

The last follow-up showed no significant changes in SNA (WMD 0.06, 95%CI:-.05, 0.16), ANB (WMD 0.19,95%CI:-0.43 ,0.82), overjet (WMD -0.53, 95%CI: -1.36, 0.30), overbite (WMD 0.20, 95%CI:-0.17, 0.57), L1/MP (WMD -0.38, 95%CI:-1.67, 0.92), while there was a significant change in SNB and U1/SN (WMD 0.38, 95%CI:0.24, 0.51) and (WMD 2.12, 95%CI:0.43, 3.82) respectively.

3.2. Subgroup analysis

A subgroup analysis based on the duration of follow-up: less than 2 years (group 1) and more than 2 years (group 2) was conducted to identify the source of the high heterogeneity and the influence of the follow-up duration.

SNA increased significantly after a 2 year-follow-up [WMD 0.07, 95%CI: 0.03, 0.11]] but no significant changes were noted in less than 2 year-follow-up [WMD -0.02, (95%CI: -0.23, 0.19)].

SNB did not have any significant changes in more than 2 year-follow-up [WMD 0.03 (95%CI: -0.11, 0.18)]; however, it increased significantly in group 1 [WMD 0.76 (95%CI: 0.49, 1.03)].

There were no significant changes in ANB in both group 2 durations [WMD .047 (95%CI:-0.26,1.20)] or group 1 [WMD -0.49 (95%CI:-1.72, 0.74)] durations. For overjet, it decreased significantly after a 2 year-follow-up [WMD -0.80 (95%CI: -1.23, -0.36)], but not in less than a 2 year-follow-up [WMD -0.49 (95%CI: -1.38, 0.41)]. Overbite grew significantly in group 2 [WMD 0.50 (95%CI:0.01, 0.98)] and not in group 1 [WMD 0.03 (95%CI: -0.36, 0.41)]. Only a few studies measured U1/SN and L1/MP; hence, the heterogeneity could not be measured due to the

small sample size. The results of the statistical testing for heterogeneity and the corresponding funnel plots are given in Figures 2c, 3c, 4c, 5c, and 6c.

3.3. Risk of bias within studies /publication bias

No publication bias was determined by using the Begg's test [in STATA 14 (StataCorp LP, College Station, USA)].

The results of the Begg's test for the analysis of small study effect (publication bias) for the measurements of SNA, SNB, ANB, overjet, overbite, U1/SN, and L1/MP are as follows, respectively: 0.78, 0.33, 0.95, 0.08, 0.45, 0.98, and 0.34.

4. DISCUSSION

4.1. Summary of evidence

This meta-analysis showed some significant relapse in skeletal and dental variables during the follow-up period. SNA and overbite increased significantly after a 2 year-follow-up. On the contrary, SNB increased significantly before a 2 year-follow-up. Overjet was significantly reduced after a 2 year-follow-up. To the best of our knowledge, this is the first meta-analysis reviewing the stability of skeletal class III malocclusion after bimaxillary surgery or mandibular setback. Data from this study revealed that the main relapse in SNA occurred after a 2 year-follow-up but not in less than 2 years. This suggested that SNA relapse often happened after a 2 year-follow-up and was largely associated with the growth of maxilla that is a common finding in class III malocclusion.

For SNB, a significant increase was noted before a 2 year-follow-up and not after a 2 year-follow-up. This relapse in the short-term is due to the growth of mandible which reportedly can continue even after 18 years of age [17]. Hence, it is important to consider the patient's age and their related growth pattern before bimaxillary surgery and/or mandibular setback treatment. Overjet was reduced significantly in more than a 2 year-follow-up; this relapse is due to an increase in SNB. However, the extent of overbite increased significantly after 2 years. Pre-surgical orthodontic treatment aims to decompensate incisor inclination toward normal values. Orthodontic decompensation allows a greater surgical correction, and this may be a more important factor in the relapse. We should keep in mind that skeletal relapse is masked frequently by compensatory changes in the axial inclination of the teeth [18-21]. Relapse varies considerably between patients and surgeons without any known reason. It is clear that good surgical training, profound experience in orthognathic surgery, and technical refinements by the surgeon are required to have perfect surgical outcomes with regards to esthetics and stability. The orthodontist should prepare the patient before surgery for a perfect coordination and leveling and alignment of both dental arches in transverse width, correct decompensation of

Table 4. Outcomes in terms of cephalometric measurements of the included studies in the quantitative meta-analysis.

Article & year	Group	Outcome			
		Before surgery (T1)	After surgery (T2)	T1-T2	Last Follow-up (T3)
Jeong et al [1], 2018	Conventional bimaxillary surgery	N/A	N/A	SNA: 0.4±2.3 SNB: -4.9±2.1	N/A
	surgery first	N/A	N/A	SNA: 0.3±2.3 SNB: -5.7±2.6	SNA: -0.1±1 SNB: 0.1±0.9
Park et al [3], 2016	Conventional bimaxillary surgery	N/A	N/A	SNA: -2.8±1.9 SNB: -3.7±1.6	N/A
	Surgery-first	N/A	N/A	SNA: -2.5±2.3 SNB: -3.2±1.9	SNA: 0.4±1.3 SNB: 0.7±1.0
Aydemir et al [4], 2015	BSSO with Semi Rigid Fixation and Lefort 1 with Rigid Fixation	SNA: 78.9±2.9 SNB: 83.1±3.9 ANB: 24.2±3.1 OJ: 27.0±3.5 OB: 22.3±2.6	SNA: 82.9±2.5 SNB: 82.4±2.9 ANB: 0.5±2.5 OJ: 2.6±3.4 OB: 1.3±1.7	SNA: 4.1±2.3 SNB: -1.4±2.6 ANB: 5.5±2.3 OJ: 9.6±3.7 OB: 3.1±3.6	SNA: 83.0±3.0 SNB: 81.4±3.6 ANB: 1.6±2.2 OJ: 3.0±1.0 OB: 1.4±0.6
					SNA: 1.1±0.3 SNB: -1±0.4 ANB: 1.1±0.1 OJ: 0.4±5.9 OB: 0.1±1.2
Kim et al [5], 2014	Bimaxillary Surgery Le Fort 1 and IVRO	SNA: 81.4±2.7 SNB: 83.8±3.7 U1/SN: 115.1±5.5 IMPA: 84.9±6.7	SNA: 81.94±2.9 SNB: 79±3.2 U1/SN: 107.9±5.4 IMPA: 83.2±5.9	SNA: 0.6±1.1 SNB: -4.9±2.4 U1/SN: -7.2±3.7 IMPA: -1.7±3.9	SNA: 0.1±0.0 SNB: 0.1±0.0 U1/SN: 0.6±0 IMPA: 1.2±1.2
	Conventional-BSSO	SNA: 81±3.8 SNB: 83.5±4.3 ANB: -2.5±2.8 IMPA: 92.6±5.9 OJ: -6.2±3.4 OB: 0.9±1.3	N/A	SNB: -4.7±1.9 IMPA: 0±0 OJ: 10.1±3.5 OB: 0.1±1.4	SNB: 0.9±0.8 IMPA: -8.9±5 OJ: -0.8±1.2 OB: 0.2±0.6
Kim et al [6], 2014	Surgery first	SNA: 80.5±4.1 SNB: 83.5±4.1 ANB: -3±2 IMPA: 82.3±9.3 OJ: -4.4±2.9 OB: 0.7±2	N/A	SNB: -4.4±1.3 IMPA: 0.7±2.2 OJ: 9.3±4.4 OB: 0.8±2.1	SNB: 1.1±0.7 IMPA: -0.6±5.7 OJ: -2.3±3.1 OB: -0.3±0.9
	Lefort 1 + BSSO	SNA: 79.6±3.6 SNB: 84.7±5.5 ANB: -5.1±4.1 OJ: -7±5 OB: -1.8±3.8	N/A	SNA: 3.3±2.1 SNB: -4±3 ANB: 7.3±3.4 OJ: 9.8±4.6 OB: 3.7±3.5	SNA: -0.1±0.9 SNB: 1±1.2 ANB: -1±1.2 OJ: -0.8±2 OB: 0±0.7

Abeltins et al[9], 2011	IVRO	SNA: 80.7±3.7 SNB: 84.9±4.9 ANB: -4.2±3.3	N/A	N/A	N/A	N/A
	BSSO	SNA: 79.3±3 SNB: 83.6±3.7 ANB: -4.2±3	N/A	N/A	N/A	N/A
Iannetti et al [10], 2007	Lefort 1	N/A	N/A	SNA: 0.1±0.1	N/A	SNA: 0.1±0.1 SNB: 0.1±0.1 ANB: 0.1±0.2
	Lefort1+BSSO	N/A	N/A	SNA: 0.1±0.1	N/A	SNA: 0.1±0.1 SNB: -0.5±0.3 ANB: 0.6±0.3
	Rigid titanium	SNA: 78.4±2.8 SNB: 81.9±3.8 ANB: -3.5±2.7 U1/SN: 111.2±7.8 IMPA: 88.3±7.1 OJ: -4.6±3.8 OB: 1.1±2.1	SNA: 81.7±2.9 SNB: 80.7±3.7 ANB: 1±2.4 U1/SN: 112.7±8.4 IMPA: 87.7±7.8 OJ: 3.2±1.31 OB: 2.3±1.4	SNA: 3.3±1.6 SNB: -2.2±2.9 ANB: 5.5±2.8 U1/SN: -2.3±3.5 IMPA: -0.2±2.7 OJ: 7.9±3.5 OB: -0.4±2.1	N/A	SNA: -0.1±0.6 SNB: 0.9±1.1 ANB: -0.9±1.3 U1/SN: 3.9±5.4 IMPA: -0.4±4.1 OJ: -0.1±1.7 OB: 1.6±1.9
Costa et al [11], 2006	Rigid resorbable plate	SNA: 80±3.6 SNB: 82.9±3.3 ANB: -2.9±0.8 U1/SN: 105.8±7.1 IMPA: 85.6±5.6 OJ: -5.1±1.9 OB: 0.8±2.1	SNA: 82.4±3.9 SNB: 80.3±3.6 ANB: 2.1±1.9 U1/SN: 107.4±7.4 IMPA: 86.1±6.5 OJ: 2±1.0 OB: 2.8±0.8	SNA: 3±1.28 SNB: -3.6±1.5 ANB: 6.6±1.4 U1/SN: 3.1±9 IMPA: 2.4±3.3 OJ: 8.2±2.4 OB: 0.9±2.4	N/A	SNA: -0.6±1.2 SNB: 1.1±0.6 ANB: -1.7±1.0 U1/SN: 1.4±3.3 IMPA: -1.9±2.9 OJ: -0.4±0.8 OB: 0.3±1.1
Kwon et al [12], 2000	Lefort 1 and BSSO	N/A	N/A	OJ: 9.1±4.6 OB: 2.5±3	N/A	OJ: -0.6±1.4 OB: -0.1±2.0
McCance et al[13], 1992	Lefort 1 +BSSO	SNA: 78.2±4.3 SNB: 83±4.4 ANB: -4.8±1.9 IMPA: 85±11 OJ: -4.1±3 OB: -6.3±3.8	SNA: 82.3±3.4 SNB: 81.4±2 ANB: 1.3±1.2 IMPA: 83.3±6.3 OJ: -0.9±1.1 OB: 2.4±1.2	SNA: 4.1±1 SNB: -2±2.0 ANB: 5.9±0.0 IMPA: -1±2.0 OJ: 2.4±4 OB: 3.2±3.6	N/A	SNA: -0.1±0.0 SNB: 0.4±1 ANB: 0.2±0.4 IMPA: -0.5±0.5 OJ: 0.8±0.0 OB: -0.7±0.5
Data are presented as means ± standard deviations OB, Over bite; OJ, Over Jet, BSSO, Bilateral sagittal split osteotomy; IVRO Intra-oral vertical ramus osteotomy, SF, Surgery-First						

the incisors, control of the surgical splint, and its newly defined occlusion to allow correct placement of the mandible during surgery. The etiology of relapse is multifactorial, including, but not limited to: the proper seating of the condyles, the extent of mandibular setback and maxillary advancement, the soft tissue and muscles, the mandibular plane angle, the remaining growth and remodeling, the skill of the surgeon, and the pre-operative age of the patient [8]. Proffit et al [22] questioned the stability in orthognathic surgery since the stability of the surgical repositioning of the jaws varies considerably depending on the procedure. In their view, the order of importance starts with the direction of movement, the type of fixation used, and in the end, the surgical technique that has been adopted.

4.2. Limitations

This meta-analysis might be considered a first step in addressing the stability of skeletal class III malocclusion after bimaxillary surgery or mandibular setback. Although this study provided an overview of the topic, there were several limitations. One main limitation was the shortage of large and high-quality RCTs. The numbers of relevant research articles and patients included in the meta-analysis were not large enough.

Furthermore, the sample sizes were diluted due to too many study variables included (7 cephalometric variables at 3 different time points). Hence, the quantitative analysis cannot accurately reflect real skeletal and dental changes. Additionally, not every study included looked at all variables further complicating the analysis.

Eventually, some studies proposed surgery first which were deleted from the analysis; however, whether the treatment effects of surgery first can be stable remains unclear. Attention should also be paid to the stability of the treatment effects of surgery first. Therefore, future research in this area is warranted.

REFERENCES

1. Chew MT, Sandham A, Wong HB. Evaluation of the linearity of soft to hard-tissue movement after orthognathic surgery. *Am J Orthod Dentofacial Orthop*. 2008;134(5):665-670. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
2. Espeland L, Hogevoeld HE, Stenvik A. A 3-year patient-centred follow-up of 516 consecutively treated orthognathic surgery patients. *Eur J Orthod*. 2008;30(1):24-30. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
3. Jakobsone G, Stenvik A, Sandvik L, Espeland L. Three-year follow-up of bimaxillary surgery to correct skeletal Class III malocclusion: stability and risk factors for relapse. *Am J Orthod Dentofacial Orthop*. 2011;139(1):80-89. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
4. Jamilian A, Cannavale R, Piancino MG, et al. Methodological quality and outcome of systematic reviews reporting on orthopaedic treatment for class III malocclusion: overview of systematic reviews. *J Orthod*. 2016;43(2):102-120. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
5. Johnston C, Burden D, Kennedy D, et al. Class III surgical-orthodontic treatment: a cephalometric study. *Am J Orthod Dentofacial Orthop*. 2006;130(3):300-309. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)

5. CONCLUSIONS

On the basis of this review, we concluded the following.

1. Surgical orthodontic improves sagittal skeletal and dental relationships but significant relapse during the follow-up period may happen.
2. SNA and overbite increased significantly after a 2 year follow-up.
3. SNB increased significantly before a 2 year follow-up with no significant changes after this follow-up.
4. Overjet diminished significantly after a 2 year follow-up.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

FOUNDING

This manuscript was partially supported by the University of Michigan Graduate Research Fund.

AUTHOR CONTRIBUTIONS

AJ: conceptualization, study design, study concept, original writing and corresponding author. LN: drafting, data interpretation, drawings and editing. MT and AD: literature review, search design, data gathering. AF and MA: software programs and statistical analysis. CWW: drafting, data interpretation, critical revision and final approval.

ACKNOWLEDGMENTS

The authors report no commercial, proprietary, or financial interest in the products or companies described in this article.

THE AUTHOR'S INSTITUTIONAL AFFILIATIONS WHERE THE WORK WAS CONDUCTED

Department of orthodontics, Faculty of dentistry, Cranio maxillofacial Research center, Tehran medical sciences, Islamic Azad University, Tehran, Iran and Department of Periodontics and Oral Medicine, School of Dentistry, University of Michigan, Michigan, USA.

6. Jamilian A, Showkatbakhsh R, Behnaz M, et al. Tooth-borne distraction osteogenesis versus conventional Le Fort I in maxillary advancement of cleft lip and palate patients. *Minerva Stomatol*. 2018;67(3):117-124. [\[Full text links\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
7. Eslami S, Faber J, Fateh A, et al. Treatment decision in adult patients with class III malocclusion: surgery versus orthodontics. *Prog Orthod*. 2018;19(1):28. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
8. Joss CU, Vassalli IM. Stability after bilateral sagittal split osteotomy setback surgery with rigid internal fixation: a systematic review. *J Oral Maxillofac Surg*. 2008;66(8):1634-1643. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
9. Iannetti G, Chimenti C, Di Paolo C. Five-year follow-up of Le Fort I osteotomies. *J Craniomaxillofac Surg*. 1987;15(5):238-243. [\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
10. LaBanc JP, Turvey T, Epker BN. Results following simultaneous mobilization of the maxilla and mandible for the correction of dentofacial deformities: analysis of 100 consecutive patients. *Oral Surg Oral Med Oral Pathol*. 1982;54(6):607-612. [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar Scopus](#)
11. Proffit WR, Phillips C, Turvey TA. Stability after surgical-orthodontic corrective of skeletal Class III malocclusion. 3.

- Combined maxillary and mandibular procedures. *Int J Adult Orthodon Orthognath Surg.* 1991;6(4):211-225.
[\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
12. de Haan IF, Ciesielski R, Nitsche T, Koos B. Evaluation of relapse after orthodontic therapy combined with orthognathic surgery in the treatment of skeletal class III. *J Orofac Orthop.* 2013;74(5):362-369.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
13. Shamseer L, Moher D, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ.* 2015;350:g7647.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#)
14. Higgins JP, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928.
[\[CrossRef\]](#) [Google Scholar](#)
15. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *J Clin Epidemiol.* 2009;62(10):e1-e34.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
16. Moher D, Liberati A, Tetzlaff J, et al. Reprint--preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Phys Ther.* 2009;89(9):873-880.
[\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
17. Love RJ, Murray JM, Mamandras AH. Facial growth in males 16 to 20 years of age. *Am J Orthod Dentofacial Orthop.* 1990;97(3):200-206.
[\[CrossRef\]](#) [Google Scholar](#)
18. Schendel SA, Epker BN. Results after mandibular advancement surgery: an analysis of 87 cases. *J Oral Surg.* 1980;38(4):265-282.
[\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
19. Vasir NS, Thompson RT, Davies TM. Dental and skeletal changes following sagittal split osteotomy for correction of mandibular prognathism. *Eur J Orthod.* 1999;13(2):134-142.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
20. Grassia V, d'Apuzzo F, DiStasio D, et al. Upper and lower arch changes after mixed palatal expansion protocol. *Eur J Paediatr Dent.* 2014;15(4):375-380.
[\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
21. Grassia V, D'Apuzzo F, Ferrulli VE, et al. Dento-skeletal effects of mixed palatal expansion evaluated by postero-anterior cephalometric analysis. *Eur J Paediatr Dent.* 2014;15(1):59-62.
[\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
22. Proffit WR, Turvey TA, Phillips C. Orthognathic surgery: a hierarchy of stability. *Int J Adult Orthodon Orthognath Surg.* 1996;11(3):191-204.
[\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
23. Jeong JH, Choi SH, Kim KD, et al. Long-term stability of pre-orthodontic orthognathic bimaxillary surgery using intraoral vertical ramus osteotomy versus conventional surgery. *J Oral Maxillofac Surg.* 2018;76(8):1753-1762.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
24. Choi SH, Cha JY, Park HS, Hwang CJ. Intraoral vertical ramus osteotomy results in good long-term mandibular stability in patients with mandibular prognathism and anterior open bite. *J Oral Maxillofac Surg.* 2016;74(4):804-810.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
25. Park KH, Sandor GK, Kim YD. Skeletal stability of surgery-first bimaxillary orthognathic surgery for skeletal class III malocclusion, using standardized criteria. *Int J Oral Maxillofac Surg.* 2016;45(1):35-40.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
26. Aydemir H, Efendiyeve R, Karasu H, Toygar-Memikoğlu U. Evaluation of long-term soft tissue changes after bimaxillary orthognathic surgery in Class III patients. *Angle Orthod.* 2015;85(4):631-637.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
27. Kim JY, Jung HD, Kim SY, et al. Postoperative stability for surgery-first approach using intraoral vertical ramus osteotomy: 12 month follow-up. *Br J Oral Maxillofac Surg.* 2014;52(6):539-544.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
28. Kim CS, Lee SC, Kyung HM, et al. Stability of mandibular setback surgery with and without presurgical orthodontics. *J Oral Maxillofac Surg.* 2014;72(4):779-787.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
29. Ko EW, Lin SC, Chen YR, Huang CS. Skeletal and dental variables related to the stability of orthognathic surgery in skeletal Class III malocclusion with a surgery-first approach. *J Oral Maxillofac Surg.* 2013;71(5):e215-e223.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
30. Abeltins A, Jakobsone G, Urtane I, Bigestans A. The stability of bilateral sagittal ramus osteotomy and vertical ramus osteotomy after bimaxillary correction of class III malocclusion. *J Craniomaxillofac Surg.* 2011;39(8):583-587.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
31. Iannetti G, Fadda MT, Marianetti TM, et al. Long-term skeletal stability after surgical correction in class III open-bite patients: A retrospective study on 40 patients treated with mono- or bimaxillary surgery. *J Craniofac Surg.* 2007;18(2):350-354.
[\[CrossRef\]](#) [Google Scholar](#) [Scopus](#)
32. Costa F, Robiony M, Zorzan E, et al. Stability of skeletal Class III malocclusion after combined maxillary and mandibular procedures: titanium versus resorbable plates and screws for maxillary fixation. *J Oral Maxillofac Surg.* 2006;64(4):642-651.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
33. Kwon TG, Mori Y, Minami K, et al. Stability of simultaneous maxillary and mandibular osteotomy for treatment of class III malocclusion: An analysis of three-dimensional cephalograms. *J Craniomaxillofac Surg.* 2000;28(5):272-277.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)
34. McCance AM, Moss JP, James DR. Stability of surgical correction of patients with Skeletal III and Skeletal II anterior open bite, with increased maxillary mandibular planes angle. *Eur J Orthod.* 1992;14(3):198-206.
[\[Full text links\]](#) [\[CrossRef\]](#) [\[PubMed\]](#) [Google Scholar](#) [Scopus](#)

Abdolreza JAMILIAN

DDS, PhD, Professor
 Department of Orthodontics
 Faculty of Dentistry
 Cranio-Maxillofacial Research Center
 Tehran Medical Sciences
 Islamic Azad University
 Tehran, Iran



CV

Professor Abdolreza Jamilian is a researcher and specialist in field of Orthodontics. He received his DDS (1991), MSc in Orthodontics (1998), and Fellowship of Orthognathic Surgery & Craniofacial Syndroms (2010) from the Shahid Beheshti University in Tehran, Iran. He obtained his European Board of Orthodontics in 2013. Now he is a professor at the Islamic Azad University in Tehran. His practice is limited to Orthodontics. He has lectured in several international congresses and has been a consultant for various journals. He has published over 200 original, peer reviewed research and review articles, 15 book chapters and more than 300 scientific communications. He holds 3 patents with the United States Patent and Trademark Office. Research interests: (1) Class 3 malocclusion (2) Cleft lip and palate (3) Orthognathic surgery.

Questions

1. Which one is correct regarding stability after surgery.

- ☐ a. Single jaw surgery has greater relapse than two jaw surgery;
- ☐ b. Two jaw surgery has greater relapse than single jaw surgery;
- ☐ c. There is a controversy regarding the stability of single and two jaw surgery;
- ☐ d. None of them.

2. What are the causes of late relapse after orthognathic surgery?

- ☐ a. Unstable occlusal relationships;
- ☐ b. Absence of myofunctional adaptation;
- ☐ c. Persistent tongue or orofacial muscle habits;
- ☐ d. All of them.

3. How much of the Class III malocclusions have true mandibular prognathism?

- ☐ a. 5% to 10%;
- ☐ b. 20% to 25%;
- ☐ c. 30% to 50%;
- ☐ d. 50% to 70%.

4. How much of the of all class III malocclusions have some degree of maxillary retrusion?

- ☐ a. 15%;
- ☐ b. 25%;
- ☐ c. 50%;
- ☐ d. 75%.



SIDO 51ST SIDO INTERNATIONAL CONGRESS
Società Italiana di Ortodonzia

AIDOR 17th AIDOR CONGRESS
Accademia Italiana DI ORTODONZIA

November 12th - 13th - 14th, 2020
Firenze, Fortezza da Basso

THE BEST OF...

- 3D – DIGITAL TECHNOLOGY
- SKELETAL ANCHORAGE
- MANAGEMENT OF THE ESTHETIC ZONE

DIGITAL PLAZA

www.sido.it